

Solutions of Maxwell Equations for Hollow curved Wave Conductor

V.Bashkov, A.Tchernomorov
Kazan State University, Kazan, RUSSIA

Abstract

In the present paper the idea is proposed to solve Maxwell equations for a curved hollow wave conductor by means of effective Riemannian space, in which the lines of motion of fotons are isotropic geodesies for a 4-dimensional space-time. The algorythm of constructing such a metric and curvature tensor components are written down explicitly. The result is in accordance with experiment.

1. Parallel translation of polarization vector in geometrical optics of non-homogenous media.

The discovery of law of parallel translation for vectors $\vec{e} = \frac{\vec{E}}{|E|}$ and $\vec{h} = \frac{\vec{H}}{|H|}$ of electric and magnific fields in media with slowly changing refraction factor is being rizen to the paper of S.M.Rytov [1] in 1938. It was shown that for the light beam of nonflat curve form there is a rotation of vectors $\vec{e}$ and $\vec{h}$ in respect to natural 3-facet, created by tangent $\vec{\tau}$, normal $\vec{n}$ and binormal $\vec{b}$ vectors to the curved beam . Setting :

$$\begin{cases} \vec{e} = \vec{n} \cos \varphi + \vec{b} \sin \varphi, \\ \vec{h} = -\vec{n} \sin \varphi + \vec{b} \cos \varphi, \end{cases} \quad (1)$$

we find

$$\frac{d\varphi}{ds} = \frac{1}{T}, \quad (2)$$

where S — arch length, T — tersion radius. Formula (2) is known as Rytov law.

In 1941 the paper of V.V.Vladimirsky [2] has appeared in which on the basis of "Rytov Law" there was predicted a global (topological) effect: the angle of rotation of polarized light beam plane, trajectory of which in optically non-homogenous media presents nonflat curve, equals integral of Gauss curvature over a region bounded by contour C , which is obtained by the top of the vector $\vec{\tau}$ on sphere of unit radius. The angle θ equals space angle π concluded inside of the conus drawn by the vector $\vec{\tau}$. This angle is coincided with topological Berry phase [3] for spiral fotons, experimentally founded in recent works of Tomita and Chao [4] in spirally curved optical wave conductors. In the case of a flat curve $\theta = 0$ and after a beam gets initial direction the field vectors take their previous position. In the first series of experiments the spiral was uniformly winded on cylinder . In the second series of experiments providing a fixed length P and radius r there were used non-uniform spirals of different shapes, modeled on computer . The space angle in this case is calculated by formula:

$$\Omega(C) = \int_0^{2\pi} \int_0^{\theta(\varphi)} \sin \theta d\theta d\varphi = \int_0^{2\pi} (1 - \cos \theta) d\varphi \quad (3)$$

Berry [3] was searching for solution of classical wave equations for a curved wave conductor by means of local coordinate system in approximation of idea of locally connected modes and weakly directing wave conductor [4]. It was produced not taking into account radiation effects, connection with reflected modes and elastic-optical effects. It led in fact to the approximation of geometrical optics and could have been analized in easier way, taking into account that light trajectory represents a broken line, corresponding to multiple reflections from walls [5],[6]. In the present paper for interpretation of observed effect we find the exact solutions of Maxwell equations for hollow curved wave conductor in certain curved Riemannian space-time different from flat Minkowsky space-time of special theory of relativity. The choice of 4-dimensional Riemannian space is such that trajectories of light beams to be isotropic geodesics in this space, so the projection of this geodesics in 3-dimensional space were the lines corresponding to topology of the wave conductor. So consider a wave conductor the axis line of which is described by equation :

$$\vec{r}_0 = \vec{i} a \cos t + \vec{j} a \sin t + \vec{k} bt \quad (4)$$

in Cartesian coordinate system . It's easy to see that curvature of this

curve is :

$$K = \frac{1}{R} = \frac{a}{a^2 + b^2}$$

and a torsion : $\chi = \frac{1}{T} = \frac{b}{a^2 + b^2}$.

Any point of wave conductor can be described by radius-vector $\vec{r} = \vec{r}_0 + \vec{\rho}$, where

$$\vec{\rho} = \eta \vec{n} + \zeta \vec{b} \quad (5)$$

$\vec{n}$ – normal vector to curve (5), $\vec{b}$ – binormal vector. The boundary of wave-conductor is described by equation $\rho = \rho_0$. Let us introduce new variable $\xi \equiv s = \sqrt{a^2 + b^2}t$ which is a length of axis line arch. Then formula for any point inside the wave conductor is :

$$\begin{cases} x = (a - \eta) \cos \frac{\xi}{\sqrt{a^2 + b^2}} - \zeta \frac{b}{\sqrt{a^2 + b^2}} \sin \frac{\xi}{\sqrt{a^2 + b^2}}, \\ y = (a - \eta) \sin \frac{\xi}{\sqrt{a^2 + b^2}} + \zeta \frac{b}{\sqrt{a^2 + b^2}} \cos \frac{\xi}{\sqrt{a^2 + b^2}}, \\ z = \frac{b\xi - a\zeta}{\sqrt{a^2 + b^2}}. \end{cases} \quad (6)$$

Let's go from coordinates (x, y, z) to locally curved coordinates (ξ, η, ζ) . The metric tensor of 3-dimensional space in this coordinate system being :

$$g_{\alpha\beta} = \begin{pmatrix} 1 + \frac{K}{a}\eta^2 + \chi^2\zeta^2 - 2K\eta & \chi\zeta & -\chi\zeta \\ \chi\zeta & 1 & 0 \\ -\chi\eta & 0 & 1 \end{pmatrix} \quad (7)$$

We suppose further that optical beams – lines of motion of fotons – are curves, which tangent vectors are orthogonal to planes $\xi = \xi_0$. It's dictated by our desire to get a non-dispersed beam at the exit of wave conductor. Congruation of these lines is described by equations

$$\begin{cases} \xi = \xi \\ \eta = r \cos(\chi\xi + \varphi) \\ \zeta = r \sin(\chi\xi + \varphi) \end{cases} \quad (8)$$

where r and φ - some parameters, characterizing the certain lines of congruation. Going to coordinates (ξ, r, φ) we get for congruation lines correspondong to optical beams :

$$\begin{cases} x = (a - r \cos(\chi\xi + \varphi)) \cos \frac{\xi}{\sqrt{a^2 + b^2}} - r \sin(\chi\xi + \varphi) \frac{b}{\sqrt{a^2 + b^2}} \sin \frac{\xi}{\sqrt{a^2 + b^2}}, \\ y = (a - r \cos(\chi\xi + \varphi)) \sin \frac{\xi}{\sqrt{a^2 + b^2}} + r \sin(\chi\xi + \varphi) \frac{b}{\sqrt{a^2 + b^2}} \cos \frac{\xi}{\sqrt{a^2 + b^2}}, \\ z = \frac{b\xi - ar \sin(\chi\xi + \varphi)}{\sqrt{a^2 + b^2}}. \end{cases} \quad (9)$$

Transition to 4-dimensional space-time with coordinates (ξ, r, φ, t) is accomplished by replacing $\overset{0}{g}_{44} = 1$ in Cartesian coordinate system in Minkowsky space on $g_{44} = -g_{11}$, so that in local coordinate system the metric tensor of 4-dimensional space-time will be:

$$g_{ij} = \begin{pmatrix} (1 - rK \cos \phi)^2 + \chi^2 r^2 & 0 & -\chi r^2 & 0 \\ 0 & 1 & 0 & 0 \\ -\chi r^2 & 0 & r^2 & 0 \\ 0 & 0 & 0 & -(1 - rK \cos \phi) \end{pmatrix} \quad (10)$$

where $\phi \equiv \chi\xi + \varphi$.

It's easy to show that congruence of isotropic geodesics of this space-time after projecting gives the lines of optical beams – trajectories of photons in 3-dimensional space, defined by formula (9). Non-vanishing components of curvature tensor of this Riemannian space-time are:

$$\begin{cases} R_{1414} = K^2(1 - rK \cos \phi) + \chi^2 K^2 r^2 - \chi^2 K^2 \cos \phi \\ R_{1424} = \chi K \sin \phi \\ R_{1434} = -\chi K r (Kr - \cos \phi) \end{cases} \quad (11)$$

Ricci tensor components:

$$\begin{cases} R_{11} = K^2 - \chi^2 \frac{K^2(\cos \phi - Kr)}{(1 - rK \cos \phi)^2} \\ R_{12} = \frac{\chi K \sin \phi}{(1 - rK \cos \phi)^2} \\ R_{13} = \chi K r \frac{(\cos \phi) - Kr}{(1 - rK \cos \phi)^2} \\ R_{44} = -K^2 - \chi^2 \frac{K^2(\cos \phi - Kr)}{(1 - rK \cos \phi)^2} \end{cases} \quad (12)$$

Scalar curvature R is:

$$R = \frac{2}{(1 - rK \cos \phi)^2} [K^2 + \chi^2 \frac{K^2(\cos \phi - Kr)}{(1 - rK \cos \phi)^2}] \quad (13)$$

And Einstein tensor components:

$$\begin{cases} G_{11} = -2\chi^2 \frac{K^2(\cos \phi - Kr)}{(1 - rK \cos \phi)^2} - \frac{\chi^2 r^2}{(1 - rK \cos \phi)^2} [K^2 + \chi^2 \frac{K^2(\cos \phi - Kr)}{(1 - rK \cos \phi)^2} Large] \\ G_{12} = \chi K \frac{\sin \phi}{(1 - rK \cos \phi)^2} \\ G_{13} = \chi \frac{K^2(\cos \phi - Kr)}{(1 - rK \cos \phi)^2} + \frac{\chi r^2}{(1 - rK \cos \phi)^2} [K^2 + \chi^2 \frac{K^2(\cos \phi - Kr)}{(1 - rK \cos \phi)^2}] \\ G_{22} = -\frac{1}{(1 - rK \cos \phi)^2} [K^2 + \chi^2 \frac{K^2(\cos \phi - Kr)}{(1 - rK \cos \phi)^2}] \\ G_{33} = -\frac{1}{(1 - rK \cos \phi)^2} [K^2 + \chi^2 \frac{K^2(\cos \phi - Kr)}{(1 - rK \cos \phi)^2}] \end{cases} \quad (14)$$

2. Solution of Maxwell equations.

In general covariant formulation the Maxwell equations are:

$$\begin{cases} F_{[ij,k]} = 0 \\ F_{,j}^{ij} = 0 \end{cases} \quad (15)$$

where comma means covariant devivative, and F_{ij} — components of electromagnetic field tensor. Writing down the system (15) in coordinates (ξ, r, φ, t) in Riemannian space (10) it can be shown that it has a partial solution corresponding to co-axial line with singularities on axis line $r = 0$ of the type :

$$\begin{cases} F_{12} = F_{24} = \frac{C_1}{r} e^{ik(\xi-t)}, \\ F_{13} = F_{34} = C_2 e^{ik(\xi-t)}, \\ F_{14} = F_{23} = 0. \end{cases} \quad (16)$$

For components of energy-momentum tensor of electromagnetic field we find:

$$\begin{cases} T_{11} = \frac{C_1^2 + C_2^2}{r^2}, \\ T_{14} = -\frac{C_1^2 + C_2^2}{r^2}, \\ T_{44} = \frac{C_1^2 + C_2^2}{r^2}. \end{cases} \quad (17)$$

Conclusion.

Returning to 3-dimensional designations $\vec{E}$ and $\vec{H}$ by means of the known formula it can be shown from (16), that along fotons lines (8) we found a vector of linear polarization rotates on the angle $\chi\xi$, what is well-coordinated with experimental data and caused by effect of topological phase of Vladimirski-Berry.

References.

1. Rytov S.M., Docl. Academ.Nauc USSR, v.28,N 4 - 5, 1938, p.263.
2. Vladimirski V.V., Docl. Academ. Nauc USSR, v.31,N 3, 1941,p.222
3. Berry M.V. Nature, 1987, v.326, N 6110, p.277
4. Tomito A.,Chaio R.V., Phis. Rev. Lett., 1986, v. 57, N8, p.937
5. Bialyniski-Birula I., Bialynski - Birula S., Phis.Rev.D., 1987, v.35, N8, p.2383
6. Vinitzki S.I., Derbov V.L., Dubrovin V.M., Marcovski B.L., Stepanovski Yu.P., Proceedings of the work conference on elaboration and construction of radiator and detector of gravitational waves, Dubna, 1989, p.74.